

AntiBak Residual • Test Results

EN 1276

Bacteria

A quantitative suspension test designed to evaluate bactericidal activity.

Pass: ≥ 5 log reduction in 5 minutes under clean conditions.

Pathogen	Pass	Dilution	Log. reduction	Test time
<i>Campylobacter jejuni</i>	✓	4%	> 5.99*	1 min.
<i>Enterococcus hirae</i>	✓	4%	> 5.25*	5 min.
<i>Escherichia coli</i>	✓	4%	> 7.00*	5 min.
<i>Klebsiella pneumoniae</i>	✓	4%	> 6.23*	1 min.
<i>Pseudomonas aeruginosa</i>	✓	20%	> 5.27*	5 min.
<i>Salmonella enteritidis</i>	✓	4%	> 6.20*	1 min.
<i>Salmonella typhimurium</i>	✓	4%	> 6.23*	5 min.
<i>Staphylococcus aureus</i>	✓	4%	> 5.27*	5 min.
Vancomycin-resistant <i>Enterococcus faecalis</i> (VRE)	✓	20%	> 6.00*	1 min.

EN 13727

Bacteria

A quantitative suspension test designed to evaluate bactericidal activity in the medical area.

Pass: ≥ 5 log reduction in 10 minutes under clean conditions.

Pathogen	Pass	Dilution	Log. reduction	Test time
<i>Legionella pneumophila</i>	✓	10%	> 5.43*	10 min.

EN 14476

Viruses

A quantitative suspension test designed to evaluate virucidal activity.

Pass: ≥ 4 log reduction in 5 minutes under clean conditions.

Pathogen	Pass	Dilution	Log. reduction	Test time
Bovine Coronavirus	✓	20%	≥ 4.40	30 sec.
Swine Flu H1N1 [†]		25%	≥ 3.25	5 min.

EN 1650

Fungi

A quantitative suspension test designed to evaluate fungicidal activity.

Pass: ≥ 4 log reduction in 15 minutes at 20°C under dirty conditions.

Pathogen	Pass	Dilution	Log. reduction	Test time
<i>Aspergillus niger</i>	✓	20%	> 5.00	15 min.
<i>Candida albicans</i>	✓	20%	> 5.00	15 min.

Residuality testing

Proven residuality up to 28 days Tested against *E.coli* and *Listeria monocytogenes* and proven to remain effective 28 days after application.
Scientific Services, Lincolnshire

Proven residuality up to 7 days Tested against MRSA, *E.coli*, *Salmonella enteritidis*, *Candida albicans*, *Klebsiella pneumoniae* and *Listeria monocytogenes* and proven to remain effective after 24 hours, 48 hours and 7 days.
Rowett Institute, Aberdeen

Proven residuality against *E.coli* Used with a steam cleaner, proven to remain highly effective against *E.coli* after 24 hours, 48 hours and 7 days after application.
Scientific Services, Lincolnshire

Independent testing

Bacteria

A quantitative suspension test designed to evaluate bactericidal activity, carried out by the Commercial Microbiology, Aberdeen.

Pathogen	Pass	Dilution	Inactivation (%)	Test time
<i>Listeria monocytogenes</i>	✓	50%	> 6.90	5 min.

Independent M.I.C. testing

Bacteria

A quantitative suspension test evaluating the Minimum Inhibitory Concentration (MIC) of AntiBak Residual against two strains of *Serratia marcescens*, carried out by the Dept of Microbiology & Biochemistry, University of the Free State, Bloemfontein, RSA.

Pass: no bacterial growth detected.

Pathogen	Pass	Dilution	Result	Test time
<i>Serratia marcescens</i> (ATCC strain)	✓	0.2%	No growth	20 min.
<i>Serratia marcescens</i> (highly resistant strain)	✓	6.3%	No growth	20 min.

* Minimum log. reduction - limited by culture † Screen testing

Manufactured by



Bio Technics Ltd.

Linton Business Park, Gourdon, Aberdeenshire, Scotland, UK DD10 0NH

Tel: +44 (0) 1561 361515 info@bio-technics.com bio-technics.com



Use biocides safely. Always read the label and product information before use. **Disclaimer:** Every effort has been made to ensure that the information presented is correct at the time of publication. However, Bio Technics Ltd does not assume and hereby disclaims any liability to any part for any loss, damage, or disruption caused by errors or omissions resulting from negligence, accident, or any other cause. The data supplied is for guidance purposes only and is therefore without guarantee.